TEST CODE **01254010**

MAY/JUNE 2019

FORM TP 2019039

EXAMINATIONS COUNCIL CARIBBEAN

CARIBBEAN SECONDARY EDUCATION CERTIFICATE® **EXAMINATION**

ADDITIONAL MATHEMATICS

Paper 01 - General Proficiency

1 hour 30 minutes

05 JUNE 2019 (a.m.)

READ THE FOLLOWING INSTRUCTIONS CAREFULLY.

- 1. This test consists of 45 items. You will have I hour and 30 minutes to answer them.
- 2. In addition to this test booklet, you should have an answer sheet.
- 3. Each item in this test has four suggested answers lettered (A), (B), (C), (D). Read each item you are about to answer and decide which choice is best.
- 4. A list of formulae is provided on page 2 of this booklet.
- 5. On your answer sheet, find the number which corresponds to your item and shade the space having the same letter as the answer you have chosen. Look at the sample item below.

Sample Item

$$\left(4^{-2}\right)^2 \div \left(\frac{1}{16}\right)^2 =$$

- (A)
- 4-1 (B)
- (C) The context to equipment
- (D)

Sample Answer








The best answer to this item is "40", so (C) has been shaded.

- If you want to change your answer, erase it completely before you fill in your new choice. 6.
- When you are told to begin, turn the page and work as quickly and as carefully as you can. 7. If you cannot answer an item, go on to the next one. You can return to that item later.
- You may use silent, non-programmable calculators to answer items. 8.

DO NOT TURN THIS PAGE UNTIL YOU ARE TOLD TO DO SO.

- The expression x 2 is a factor of 1.
- Which of the following graphs BEST represents f(x) = x(1-x)?


- $4x^4 2x^2 56$ (A)
- $4x^3 + 2x^2 16$ (B)
- $2x^3 + 2x^2 4x 8$ (C)
- $3x^4 10x^3 5x^2 + 4$ (D)




- When $x^3 7x^2 + 2x 1$ is divided by x + 2, 2. the quotient is
 - (A) -17
 - -39(B)

484


- $x^2 5x 8$ (C)
- $x^2 9x + 20$ (D)



Given that $f(x) = ax^2 + bx + c$, f(x) can be 3.



(A) $a\left(x+\frac{b}{a}\right)^2 = \frac{ac-b^2}{a^2} = \frac{ac-b^2}{a^2}$



- $a\left(x+\frac{b}{2a}\right)^2+\frac{ac-b^2}{a^2}$ the desired and a substant with the contraction of th
- (C) $wa \left(x + \frac{b}{2a}\right)^2 + \frac{4ac b^2}{4a}$ is some across and a prediction of the values of x for which $(x + 15)^2 = 64$ The values of x for which $(x + 15)^2 = 64x$ are
- $a\left(x+\frac{b}{2a}\right)^2+\frac{4ac-b^2}{4a^2}$

- 3 and 5 (A)
- 9 and 5 (B)
- (C) 3 and 25
- 9 and 25 (D)



(A) (B) (B) (D)

use it completely before you fill in your new choice.

- The range of values for which 6. $x^2 - 7x + 10 < 0$ is
- The best answer to $t_0 < x < 2$ (A) (C) has been shaded.
 - If you want to cham 6 > x > 2, (B)
 - x < 2 and x > 5(C)
- When you 2=< x bns 2> x arm (**O**) page and work as quarkly und as carefully as you can. If you cannot answer an item, go on to the next end. You can reugh to that dain later.

nou may use silent non-programm this calculators to asswer item, c

7. The set of values of x for which $\frac{2x+1}{x-1} \ge 0$

If function $m: x \to 5 + 2x$, then m(4-2a)10.

$$(A) x > 1$$

(A)
$$4-4a$$

(B) $9-2a$

$$(B) x \ge -\frac{1}{2}$$

(C)
$$8-4a$$
 (D) $13-4a$

(C)
$$x \ge -\frac{1}{2}$$
 and $x \ge 1$

11. If
$$f^{-1}: x \to x^2 - 1, x \ge 0$$
, then

(D)
$$x \le \frac{1}{2} \text{ or } x > 1$$

(A)
$$f: x \to 1-x^2, x \in R$$

(D)
$$x \le \frac{1}{2} \text{ or } x > 1$$

(C)
$$f: x \to \sqrt{x-1}, x \ge 1$$

(B) $f: x \to \sqrt{x+1}, x \ge -1$

8. If
$$f(x) = -\frac{2}{9}x^3$$
, $-3 \le x \le 3$, $x \in R$, then

(D)
$$f: x \to \frac{2}{x^2 - 1}, x \neq \pm 1$$

$$(A) 0 \le f(x) \le 6$$

(B)
$$-6 \le f(x) \le 0$$

$$(C) -6 \le f(x) \le 6$$

$$\frac{2^{-1}}{8^{\frac{1}{3}}}$$
 simplifies to

(D)
$$6 \le f(x) \le -6$$

(A)
$$\frac{1}{4}$$

The functions f and g are defined as 9. follows:

(B)
$$\frac{1}{2}$$

$$f: x \to \frac{x+1}{x-1}, x \ne 1, x \in R$$

(c)
$$\sqrt{2}$$
 or the subsystem of f

$$g: x \to 2x + 1, x \neq \frac{1}{2}, x \in R$$

(D)
$$\frac{1}{\sqrt{2}}$$
 $\frac{1}{2}$ (A)

The function fg(x) is given by

$$(A) \qquad \frac{x+1}{x}, \ x \neq 0, \ x \in R$$

(B)
$$\frac{x-2}{x+1}$$
, $x \neq -1$, $x \in R$

(C)
$$\frac{x+1}{x+2}$$
, $x \neq -2$, $x \in R$

(D)
$$\frac{x-1}{2x+1}$$
, $x \neq -\frac{1}{2}$, $x \in R$

- 13. Given that $\log_{\rho} X = 6$ and $\log_{\rho} Y = 4$, the value of $\log_{\rho} \left(\frac{X}{Y} \right)$ is
 - (A) 10
 - (B) $\log_{p} 2$
 - (C) $\frac{\log_p 6}{\log_p 4}$
 - (D) 2
- Given that $3 \times 27^{2x} = 9^x$, the value of x is
 - (A) $-\frac{1}{4}$
 - (B) -I
 - (C) $\frac{1}{4}$
 - (D)
 - 15. The value of x for which $4^{x+1} = 2$ is

(U)

- (A) $-\frac{1}{2}$
- (B) 0

A006

- (C) $\frac{1}{2}$
- (D) 1
- 16. The value of x such that $\log_2 (5x + 1) \log_2 (3x 5) = 2$ is
 - (A) 2
 - (B) 3
 - (C) 5
 - (D) HI U 2000

- 17. The series $-2 + \frac{4}{3} \frac{8}{9} + \dots$ converges to the limit
 - (A) $-\frac{6}{5} 6 \times (81)$
 - $(B) \frac{6}{5} \le 1$
 - (C) -6
 - (D) 6
- 18. The sum of the first n terms of a series is given by $\sum_{r=1}^{n} (5-3r)$. The sum of the first 10 terms is
 - (A) -170
 - (B) -125
 - (C) 6-115() 20 (U)
 - (D) -85
- The sum of $\sum_{k=1}^{3} \frac{1}{k}$ is 19.
 - (A) $\frac{1}{3}$
 - $\begin{array}{cccc}
 \text{(B)} & \frac{1}{2} & & & \text{(A)} \\
 \text{(B)} & \frac{1}{2} & & & & \text{(A)}
 \end{array}$
 - (C) $\frac{3}{5}$ $\frac{5-7}{1-3}$ (8)
 - (D) $\frac{11}{6}$ (O)

- (A) 200
- (B) $\frac{500}{3}$
- (C) 300
- (D) $\frac{2500}{3}$

21. A line L passes through the point (6, 5) and is perpendicular to a line whose equation is 3x + 4y - 7 = 0. The equation of L is

- (A) 4x 3y 9 = 0
- (B) 4x + 3y 7 = 9
- (C) 3x + 4y 11 = 0
- (D) 3x-4y-30=0

22. The lines 7x - 4y + 25 = 0 and 3x - y - 5 = 0 intersect at the point P. The coordinates of P are

- (A) (5, 10)
- (B) (-1, 8)
- (C) (9, 22)
- (D) (-9, -32)

23. A circle C has centre (3, -2) and radius 4. The equation of C is

- (A) $x^2 + y^2 3 = 0$
- (B) $x^2 + y^2 + 6x 4y + 3 = 0$
- (C) $x^2 + y^2 6x + 4y 3 = 0$
- (D) $x^2 + y^2 + 3x 2y 3 = 0$

24. Two vectors are equal if they

- (A) have the same magnitude and different directions
- (B) have the same magnitude and same direction
- (C) are parallel and are in different directions
- (D) have different magnitudes and are in the same direction

25. The triangle OAB has vertices given by

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} -3 \\ a \end{pmatrix}$$
 and $\begin{pmatrix} 6 \\ 2 \end{pmatrix}$ respectively. Given

that the angle AÔB is $\frac{\pi}{2}$, then a =

- (A) 18
- (B) 9
- (C) -9
- (D) 0

26. Given that $OA = \begin{pmatrix} -17 \\ 25 \end{pmatrix}$ and $OB = \begin{pmatrix} 4 \\ 5 \end{pmatrix}$, then the vector $AB = \begin{pmatrix} 4 \\ 5 \end{pmatrix}$

- $(A) \qquad \begin{pmatrix} -13 \\ 30 \end{pmatrix}$
- (B) $\begin{pmatrix} -13 \\ -20 \end{pmatrix}$

(C) $\begin{pmatrix} -21 \\ 20 \end{pmatrix}$

X. aucusures

(D) $\begin{pmatrix} 21 \\ -20 \end{pmatrix}$

27. Which of the following formulae are used to calculate the area of a sector, A, with radius, r, arc length, l, and angle, q, in radians?


$$1. \quad A = \frac{r^2q}{2}$$

II.
$$A = \frac{l^2}{2q}$$

III.
$$A = \frac{lr}{2}$$

- (A) I and II only
 - (B) I and III only
 - (C) II and III only
 - (D) I, II and III

<u>Item 28</u> refers to the following triangle (not drawn to scale).





28. The size of the missing angle, X, measured in radians, is


(1)


- (A) $\frac{\pi}{5}$
- **(B)** $\frac{\pi}{10}$ $\frac{\pi}{0}$ **(a)**
- (C) $\frac{\pi}{20}$
- (D) $\frac{\pi}{25}$

29. Which of the following graphs represents $y = \sin x$?









- 30. For $0^{\circ} < x < 360^{\circ}$, the equation $3 \sin^2 x = 8 \cos x$ has only one real solution given by
 - $(A) \quad \sin x = -\frac{1}{3}$
 - (B) $\cos x = \frac{1}{3}$
 - (C) $\sin x = \frac{1}{3}$
 - (D) $\cos x = -\frac{1}{3}$

31. $\sin (\alpha + 45^{\circ})$ is equal to

(A)
$$\frac{1}{\sqrt{2}}(\sin\alpha + \cos\alpha)$$

(B)
$$\frac{1}{\sqrt{2}}(\cos\alpha - \sin\alpha)$$

(C)
$$\frac{1}{2}(\sin\alpha - \cos\alpha)$$

(D)
$$\frac{1}{2}(\cos\alpha - \sin\alpha)$$

32. $\tan\left(2x+\frac{\pi}{4}\right)$ is equal to

484

A006

(A)
$$\tan 2x + 1$$

(B)
$$\frac{\tan 2x - 1}{1 + \tan 2x}$$

$$(C) \qquad \frac{\tan 2x + 1}{1 - \tan 2x}$$

$$(D) \qquad \frac{2 \tan 2x}{1 - \tan^2 2x}$$

33. If $\sin (x + 20^\circ) = \cos x$, then the value of x is

- (A) 35°
- (B) 45°
- (C) 55°
- (D) 70°

$$\int (\sin x - \cos x) dx$$
(A). $2 \sin x - \cos x^2$
(B) $-\cos x - 2 \sin x$
(C) $\cos x - 2 \sin x^2$
(L1) $-\cos x - \sin x^2$

34. For $0 \le \theta \le \frac{\pi}{2}$, $\tan \theta = \frac{4}{3}$. The exact

value of tan 2θ is

- (A) $\frac{8}{3}$
- (B) $\frac{24}{25}$
- (C) $-\frac{24}{7}$
- (D) $-\frac{12}{7}$

35. $\frac{d}{dx}\sqrt{(7x^2+4)} =$

- $(A) \qquad \frac{14x}{\sqrt{7x^2+4}}$
- $(B) \qquad \frac{7x}{\sqrt{7x^2+4}}$
- $(C) \qquad \frac{7x}{2\sqrt{7x^2+4}}$

 $\frac{\text{over and to } \frac{ds}{ds}}{(D)^{3/s}} = \frac{\text{noise will include } \sqrt{3}T}{\sqrt{7x^2!} + 4} \times 10^{-s} \text{ on } s = 1$

36. At the point (7, 4) on the curve y = f(x),

 $\frac{dy}{dx} = 0$ and $\frac{d^2y}{dx^2} = -5$. The point (7, 4) is

((1)

- (A) an optimum point
- (B) a point of inflexion
- (C) a minimum turning point
- (D) a maximum turning point

40.

(B)
$$-\frac{1}{2}$$

(C)
$$\frac{1}{2}$$

(D)
$$\frac{\sqrt{3}}{2}$$

184

Given that $y = \cos 2x$, then $\frac{dy}{dx} = \cos 2x$ 38.

(A)
$$\sin 2x$$

(C)
$$-2 \sin 2x$$

(D)
$$-\frac{1}{2}\sin 2x$$

The gradient function $\frac{dy}{dy}$ of the curve 39. $y = \sin(2x^2 + 1)$ is

(B)
$$\cos(2x^2+1)$$

$$\frac{1}{4x} \frac{(1-x^2) \operatorname{miod} (1-x^2) \cdot (1-x^2$$

(D)
$$-4x \cos(2x^2 + 1)$$

Inion gainess mussinings (1)

noisonal remogra

The region bounded by the curve $y = x^2$, the x-axis and the lines x = 0 and x = 1 is rotated 360° about the x-axis. The volume of the solid generated can be found from

$$(A) \qquad \pi \int_0^1 x^2 dx$$

(B)
$$(a_1) \int_0^1 x^2 dx$$
 (C)

(C)
$$\int_0^1 x^4 dx$$

(D)
$$\pi \int_0^1 x^4 dx$$

 $\int (2x-5)^3 dx =$ 41.

or so then the value of x

(A) $\frac{(2x-5)^4}{4} + c$

 $\frac{(2x-5)^2}{2}+c$ (B)

 $\frac{(2x-5)^4}{8}+c$

(D)
$$\frac{2(2x-5)}{4} + c$$
 (8)

 $\int (\sin x - \cos x) dx$ 42.

(A)
$$2 \sin x - \cos x + C$$

(B)
$$-\cos x - 2\sin x + C$$

(C)
$$\cos x - 2 \sin x + C$$

(D)
$$-\cos x - \sin x + C$$

184

If $\int_{2}^{a} (6+3x)dx = 72$ where a > 2, then a =

- (A)
- (B) 10

6

- (C) 36
- (D) 72

44. If
$$X = \int_{h}^{a} f(x)dx$$
 and $a < c < b$, then $X =$

(A)
$$\int_0^a f(x)dx + \int_0^b f(x)dx$$

(B)
$$\int_a^c f(x)dx + \int_b^c f(x)dx$$

(C)
$$\int_a^c f(x)dx + \int_c^b f(x)dx$$

(D)
$$\int_0^a f(x)dx + \int_0^c f(x)dx - \int_0^b f(x)dx$$

- The region R is enclosed by the x-axis, the curve $y = x^2 + 2x 1$, the lines x = 2 and x = 3. The 45. area of R in units2 is
 - (A) 15
 - (B)
 - (C)
 - (D)

END OF TEST

IF YOU FINISH BEFORE TIME IS CALLED, CHECK YOUR WORK ON THIS TEST.

01254010/MJ/CSEC 2019